

DYNAMICS SEMI-CONJUGATED TO A SUBSHIFT FOR SOME POLYNOMIAL MAPPINGS IN \mathbb{C}^2

GABRIEL VIGNY

Abstract

We study the dynamics near infinity of polynomial mappings f in \mathbb{C}^2 . We assume that f has indeterminacy points and is non constant on the line at infinity L_∞ . If L_∞ is f -attracting, we decompose the Green current along itineraries defined by the indeterminacy points and their preimages. The symbolic dynamics that arises is a subshift on an infinite alphabet.

1. Introduction

We are interested in the dynamics of polynomial mappings f in \mathbb{C}^2 whose meromorphic extensions to \mathbb{P}^2 admit indeterminacy points and for which the line at infinity (which we denote by L_∞) is f -attracting (that is: there exists $C > 1$ such that for $p \in \mathbb{C}^2$ with $\|p\|$ large enough, one has $\|f(p)\| \geq C\|p\|$). In particular, given any large ball \mathbb{B} in \mathbb{C}^2 , these maps are polynomial-like in the sense of [DS03] from $f^{-1}(\mathbb{B})$ to \mathbb{B} . The dynamics is studied there: there exists an invariant probability measure which is K-mixing and of maximal entropy. Our goal is to study the dynamics near infinity, especially the structure of the *Green current*, which is a positive closed current of bidegree $(1, 1)$ invariant under the action of f^* .

In [DDS05], the authors consider the case where f_∞ , the restriction of f to L_∞ , is constant and they decompose the Green current into pieces associated to an itinerary defined by indeterminacy points. On the basin of attraction of the indeterminacy set, the itinerary map semi-conjugates f to a shift.

Another case which has been studied is when f admits a holomorphic extension to \mathbb{P}^2 : in [BJ00], the authors showed the Green current admits a local laminar decomposition consisting of the stable manifolds of f at

2000 *Mathematics Subject Classification.* 37F10, 37B10.

Key words. Polynomial mapping, Green current, subshift.

the points of the Julia set of f_∞ . Applying one dimensional theory, one also obtains in this case a dynamics semi-conjugated to a shift.

We study here a mixed situation. We assume that f admits indeterminacy points on L_∞ and that f_∞ is not a constant function. In order to describe clearly the new phenomena happening here, we consider the case where f_∞ is hyperbolic. The method we use allows to study more general cases. We will complete our study by giving several examples. In the hyperbolic case, we show that the Green current decomposes along some itineraries defined by the indeterminacy points and their preimages. Surprisingly, the local stable manifolds associated to the points of the Julia set of f_∞ are not charged by the Green current. Furthermore, the symbolic dynamics we obtain is a subshift (a Markov chain), which is new for polynomial mappings.

The main tools we use are horizontal-like maps and a theorem of convergence of currents proved in [Duj04] and [DDS05]. Roughly speaking, such applications are contracting in the vertical direction and expanding in the horizontal one in some bidisk. For the reader's convenience, we give the basic properties of these objects.

Next, we define and study the basic properties of the family \mathcal{G} of maps we consider. We give a simple sufficient condition for a map f to be in \mathcal{G} and we prove the algebraic stability. Then, by a theorem of Sibony [Sib99], one can associate to f a natural invariant current (Green current). We give an easily computable formula for the trace of the Green current at infinity. This trace is a probability measure which is a combination of Dirac masses at the indeterminacy points and their preimages. Under some additional hypothesis, we also compute the topological degree.

We then study the decomposition of the Green current on a neighborhood of infinity under the hypotheses that the indeterminacy set is located in the Fatou set of f_∞ , with no indeterminacy point being periodic for f_∞ and that f_∞ is hyperbolic. This set of maps contains an open subset of \mathcal{G} . The decomposition of the Green current semi-conjugates f to a subshift on an infinite alphabet. A classic object in this setting is the escape rate which measures the asymptotic speed at which a point goes to infinity (see [DS04] and [FJ05] for some interesting examples and results on this topic). Under some additional hypothesis, we show that the range of the escape rate is a full interval which is new for polynomial maps and we compute a mean escape rate. We will explain briefly how to obtain a weaker decomposition of the Green current in a more general case. Finally, we study examples, in particular the case where the indeterminacy points are located in the exceptional set of f_∞ , in

this case the support of the Green current is strictly contained in the Julia set of f .

2. Polynomial maps with dynamics at infinity

2.1. Horizontal-like maps.

We recall here the facts we use on horizontal-like maps. Proofs and details can be found in [Duj04] and [DDS05].

Let \mathbb{D} (resp. \mathbb{D}_r) be the unit disk (resp. the disk of radius r centered at 0) in \mathbb{C} . Let Δ be the unit bidisk in \mathbb{C}^2 , we denote its vertical boundary by $\partial_v\Delta$, and its horizontal boundary by $\partial_h\Delta$. Namely:

$$\partial_v\Delta = \{(z, w) \in \mathbb{C}^2, |z| = 1, |w| < 1\} \quad \text{and}$$

$$\partial_h\Delta = \{(z, w) \in \mathbb{C}^2, |z| < 1, |w| = 1\}.$$

We have the following definitions:

Definition 2.1. Let $\Delta_i \subset M_i$ be an open subset biholomorphic to Δ in the complex surface M_i for $i = 1, 2$. Let f be a dominating meromorphic map defined in some neighborhood of Δ_1 with values in M_2 . The triple (f, Δ_1, Δ_2) defines a horizontal-like map if:

- f has no indeterminacy points in $\partial_v\Delta_1$ and $f(\partial_v\Delta_1) \cap \overline{\Delta_2} = \emptyset$;
- $f(\overline{\Delta_1}) \cap \partial\Delta_2 \subset \partial_v\Delta_2$;
- $f(\Delta_1) \cap \Delta_2 \neq \emptyset$.

Definition 2.2. A positive closed $(1, 1)$ -current T in Δ is *vertical* if:

$$\text{Supp } T \subset \mathbb{D}_{1-\varepsilon} \times \mathbb{D} \text{ for some } \varepsilon > 0.$$

Similarly, we can define horizontal currents.

We can define the (horizontal) slice measures m^{w_0} of a vertical positive closed $(1, 1)$ -current T by $T \wedge [w = w_0]$. These measures have the same mass, which we call the *slice mass* of T . The current T is zero if and only if its slice mass is zero. The main fact is that we can define the pull-back of such a current by a horizontal-like map, and we have the following:

Let (f, Δ_1, Δ_2) be a horizontal-like map then there exists a positive integer $d \geq 1$ such that for every vertical positive closed current T in Δ_2 of slice mass 1, $\frac{1}{d}f^(T)$ is a vertical positive closed current in Δ_1 of slice mass 1.*

We call this integer the degree of f , it can be computed as the number of intersections of the preimage of a vertical line with a horizontal line

(with multiplicity). The following result is our main tool to obtain the convergence in Theorem 3.5:

Theorem 2.3 ([DDS05]). *Let $\{(f_i, \Delta_i, \Delta_{i+1})\}_{i \geq 1}$ be a sequence of horizontal-like maps of degree d_i such that $(f_i)^{-1}(\Delta_{i+1}) \subset \mathbb{D}_{1-\varepsilon} \times \mathbb{D} \subset \Delta_i$ for a fixed $\varepsilon > 0$. Assume that $K = \bigcap_{n \geq 1} f_1^{-1} \dots f_n^{-1}(\Delta_{n+1})$ has zero Lebesgue measure. For each n , let T_n be a vertical positive closed $(1, 1)$ -current of slice mass 1 in Δ_n .*

Then, the sequence of iterated pull-back $\left(\frac{1}{d_1} f_1^ \dots \frac{1}{d_n} f_n^* T_{n+1}\right)_n$ converges to a vertical positive closed current τ of slice mass 1 in Δ_1 which is independent of (T_n) .*

2.2. The class \mathcal{G} .

We say that the line L_∞ is attractive for a polynomial mapping f of \mathbb{C}^2 if there are constants $C > 1$ and M large enough such that for $\|p\| \geq M$, we have $\|f(p)\| \geq C\|p\|$. We can consider the meromorphic extension of f to \mathbb{P}^2 , which can have indeterminacy points, we still denote by f that extension and by $I(f)$ the indeterminacy set. Let f_∞ be the unique map extending $f|_{L_\infty \setminus I(f)}$. The case where f_∞ is constant was studied in [DDS05], so we will consider the following set of mappings \mathcal{G} :

Definition 2.4. Let \mathcal{G} the set of mappings f satisfying the following properties:

- The line L_∞ is attractive.
- The meromorphic extension of f to \mathbb{P}^2 admits indeterminacy points.
- The map f_∞ is not constant.

Let $f = (f_1, f_2)$ of algebraic degree D be in \mathcal{G} . We denote by f_1^+ and f_2^+ the homogeneous parts of maximal degree. After a linear change of coordinates, we can assume $\deg f_1^+ = D$ and $\deg f_2^+ = D' \leq D$. The meromorphic extension of f to \mathbb{P}^2 is given by $[T^D f_1(Z/T, W/T) : T^D f_2(Z/T, W/T) : T^D]$ and the restriction of f to $L_\infty = (T = 0)$ is $f_\infty[Z : W] = [f_1^+(Z, W) : 0^{D-D'} f_2^+(Z, W)]$. Thus, in order to have f_∞ not constant, we need $D = D'$ and f_1^+ not proportional to f_2^+ (otherwise, f sends L_∞ to $[1 : 0 : 0]$ or $[1 : \lambda : 0]$).

The indeterminacy set $I(f)$ of f is the common zeros of f_1^+ and f_2^+ : if the line \mathcal{D} of equation $a_j z - b_j w = 0$ satisfies $f_1^+(\mathcal{D}) = \{0\}$ and $f_2^+(\mathcal{D}) = \{0\}$ then $[b_j : a_j : 0]$ is in $I(f)$.

One deduces from above that all the mappings of \mathcal{G} can be written as:

$$(1) \quad f(z, w) = \left(\prod_{j \leq m} (a_j z - b_j w)^{\alpha_j} P_1(z, w) + Q_1(z, w), \right. \\ \left. \prod_{j \leq m} (a_j z - b_j w)^{\alpha_j} P_2(z, w) + Q_2(z, w) \right),$$

where the a_j and b_j are complex numbers satisfying $(a_j, b_j) \neq (0, 0)$, m and the α_j are positive integers, P_1 and P_2 are homogeneous polynomials of degree $d' \geq 1$ with no common factor and the Q_j are polynomials of degree strictly smaller than the degree of f . We denote by d the sum $\sum_{j \leq m} \alpha_j$, so that f has degree $d + d' = D$.

So we have that $f_\infty([Z : W]) = [P_1(Z, W) : P_2(Z, W)]$. We define the multiplicity of an indeterminacy point I as the intersection multiplicity at I of L_∞ and $f^{-1}(L)$ where L is a generic line. The indeterminacy points of f are the $I_j = [b_j : a_j : 0]$ with multiplicity α_j . We assume of course that the (a_j, b_j) are not proportional.

The following proposition shows that we can find f with any given set of indeterminacy points with multiplicity and any given restriction at infinity. Furthermore, it shows that for $D \geq 3$, \mathcal{G} corresponds to a Zariski open set of the space of parameters of (1).

Proposition 2.5. *Let $f = (f_1, f_2)$ be as in (1). Assume that the polynomial $\Phi = f_1 P_2 - f_2 P_1$ has degree $\geq 2 + d'$. If for all j , $a_j z - b_j w$ does not divide the homogeneous part of maximal degree of Φ , then L_∞ is f -attracting.*

Proof: Let f be as above and N be a small neighborhood of infinity. Observe that for any neighborhood V of $I(f)$, there exists a constant C such that if $p = (z, w) \in N \setminus V$ we have $C \|p\|^D \leq \|f(p)\|$. So we just have to prove the estimate on V . Since P_1 and P_2 have no common factor, there is $\lambda > 0$ such that $\max(|P_1(z, w)|, |P_2(z, w)|) \leq \lambda \|(z, w)\|^{d'}$ on N . The hypothesis implies that $|\Phi(z, w)| \gtrsim \|(z, w)\|^{\deg \Phi}$ near $I(f)$, hence:

$$2\|f(z, w)\| \geq \frac{|\Phi(z, w)|}{\max(|P_1(z, w)|, |P_2(z, w)|)} \gtrsim \|(z, w)\|^2.$$

The proposition follows. □

Observe that for a generic map $g \in \mathcal{G}$, we have $\deg \Phi = 2d' + d - 1$. The criterion is not optimal, but it is generic for $D \geq 3$ and easy to check. If $D = 2$, we obtain in the same way that $\|f(z, w)\| \gtrsim \|(z, w)\|$,

so we may have to multiply f satisfying the above criterion by a large enough constant in order to have that L_∞ is attractive.

Recall that a meromorphic mapping $f: \mathbb{P}^2 \rightarrow \mathbb{P}^2$ is said to be *algebraically stable* if no algebraic curve is sent to an indeterminacy point after some iterations, equivalently, if f is of algebraic degree D then f^n has degree D^n for all $n \geq 1$. It is clear that the mappings of \mathcal{G} are algebraically stable because no algebraic curve can be sent on an indeterminacy point. So by [Sib99], we can define the *Green current* for a map f in \mathcal{G} (see the discussion before Proposition 2.8 for details).

We use the notation of (1) in the following proposition.

Proposition 2.6. *Let f and g be in \mathcal{G} then $f \circ g \in \mathcal{G}$. More precisely, if $f = (PQ_1 + R_1, PQ_2 + R_2)$ has degree D and $g = (P'Q'_1 + R'_1, P'Q'_2 + R'_2)$ has degree D' then $f \circ g = (P''Q''_1 + R''_1, P''Q''_2 + R''_2)$ where:*

$$P'' = (P')^D P(Q'_1, Q'_2), \quad Q''_1 = Q_1(Q'_1, Q'_2), \quad \text{and} \quad Q''_2 = Q_2(Q'_1, Q'_2).$$

In particular, $f \circ g$ has degree $D + D'$ and for $n \in \mathbb{N}^$ we have $(f_\infty)^n = (f^n)_\infty$.*

Proof: With the above notations, the homogeneous part of maximal degree of the components of $f \circ g$ are equal to:

$$P(P'Q'_1, P'Q'_2)Q_1(P'Q'_1, P'Q'_2) = (P')^D P(Q'_1, Q'_2)Q_1(Q'_1, Q'_2)$$

and

$$P(P'Q'_1, P'Q'_2)Q_2(P'Q'_1, P'Q'_2) = (P')^D P(Q'_1, Q'_2)Q_2(Q'_1, Q'_2).$$

We only have to check that $Q_1(Q'_1, Q'_2)$ and $Q_2(Q'_1, Q'_2)$ have no common factor: if not, since two homogeneous polynomials have no common factor if and only if they have no common non trivial zero and since Q_1 and Q_2 have no common factor, we would have that Q'_1 and Q'_2 have a non trivial common zero. \square

2.3. Multiplicity of the indeterminacy points, trace of the Green current at infinity.

Let E denote the set $\bigcup_{n \geq 0} f^{-n}(I(f)) = \bigcup_{n \geq 0} I(f^n)$. For $p \in E$, we denote by $\lambda_{p,n}$ the real number equal to the multiplicity at p of f^n as an indeterminacy point divided by D^n , that is: $\lambda_{p,n} = \frac{\text{mult}_p(f^n)}{D^n}$ (these numbers will appear in the symbolic dynamics of f). We have the following lemma:

Lemma 2.7. *For all $p \in E$, $(\lambda_{p,n})$ is an increasing sequence bounded by 1. Let λ_p be its limit. Then:*

$$\sum_{p \in E} \lambda_p = 1.$$

Proof: Write $f^n = (P_n Q_{1,n} + R_{1,n}, P_n Q_{2,n} + R_{2,n})$. Recall that $I(f^n)$ is the intersection of L_∞ with the zero set of P_n . By Proposition 2.6:

$$P_{n+1} = (P_n)^D P(Q_{1,n}, Q_{2,n}).$$

Hence, $(\lambda_{p,n})$ is increasing since $(P_n)^D$ is a factor of P_{n+1} .

Set $d_n = \deg(P_n)$ and $d'_n = \deg(Q_{i,n})$. We deduce from Proposition 2.6:

$$d'_n = (d')^n \quad \text{and} \quad d_n = D^n - (d')^n.$$

So, $\sum_{p \in E} \lambda_{p,n} = \frac{d_n}{D^n} \rightarrow 1$. This completes the proof. □

Remarks.

1. In a way, the indeterminacy points of f^n take asymptotically all the available degree, so they carry the main part of the dynamics near L_∞ (cf. Proposition 2.8).
2. The sequence $(\lambda_{p,n})_n$ can be strictly increasing as we will see in the last example of Section 3.6. One can check that $(\lambda_{p,n})_n$ is strictly increasing after some rank if and only if p is preperiodic.

Recall that, on \mathbb{C}^2 , for f algebraically stable of degree D , the sequence of positive functions $(u_n = \frac{1}{D^n} \log^+ \|f^n(z, w)\|)$ almost decreases (i.e. $(u_n + c_n)_n$ is decreasing for some sequence of constant $(c_n)_n$ decreasing to zero) to the *Green function* u of f which is a potential of the *Green current* T of f ($dd^c u = T$). Furthermore, the function $\tilde{u}(z, w) = u(z, w) - \frac{1}{2} \log(|z|^2 + |w|^2 + 1)$ is a bounded quasi-plurisubharmonic function on \mathbb{C}^2 , thus it extends to \mathbb{P}^2 , and this extension satisfies $dd^c \tilde{u} = T - \omega_{FS}$ where ω_{FS} is the Fubini-Study form on \mathbb{P}^2 (see [Sib99]).

We will see in Proposition 2.8 that $\tilde{u}|_{L_\infty}$ is not identically equal to $-\infty$ so we can define the measure $m_\infty = T \wedge [L_\infty]$ which is the trace of the Green current at infinity. Since the sequence of functions $\tilde{u}_n(z, w) := u_n(z, w) - \frac{1}{2} \log(|z|^2 + |w|^2 + 1)$ is almost decreasing, m_∞ is the limit in the sense of current of the sequence $((dd^c \tilde{u}_n(z, w) + \omega_{FS}) \wedge [L_\infty])$. In particular, we have $m_\infty = dd^c(\tilde{u}|_{L_\infty}) + (\omega_{FS})|_{L_\infty}$. The next proposition shows that m_∞ is a combination of Dirac masses at the points of E , with computable coefficients.

Proposition 2.8. *Let f be in \mathcal{G} and \tilde{u} be as above. For $p \in E$, we denote by $[a_p : b_p : 0]$ its homogeneous coordinates. Then:*

$$\tilde{u}([z : w : 0]) = \log \left(\prod_{p \in E} |a_p w - b_p z|^{\lambda_p} \right) - \frac{1}{2} \log(|z|^2 + |w|^2).$$

In particular, we have the formula:

$$m_\infty = \sum_{p \in E} \lambda_p \delta_p$$

where δ_p is the Dirac mass at p .

Proof: With the above notations, we have that in \mathbb{C}^2 :

$$\begin{aligned} \tilde{u}_n(z, w) &= \frac{1}{D^n} \log^+ \|(P_n Q_{1,n} + R_{1,n})(z, w), (P_n Q_{2,n} + R_{2,n})(z, w)\| \\ &\quad - \frac{1}{2} \log(|z|^2 + |w|^2 + 1). \end{aligned}$$

So, first outside of E , and hence everywhere on L_∞ by semi-continuity, the extension is given by:

$$\begin{aligned} \tilde{u}_n([z : w : 0]) &= \frac{1}{D^n} \log \|(P_n Q_{1,n})(z, w), (P_n Q_{2,n})(z, w)\| \\ &\quad - \frac{1}{2} \log(|z|^2 + |w|^2). \end{aligned}$$

By definition of the $\lambda_{p,n}$, there is a constant C_n depending on the choice of the coordinates of the elements of E such that:

$$\begin{aligned} \tilde{u}_n([z : w : 0]) &= \sum_{p \in E} \lambda_{p,n} \log |a_p w - b_p z| \\ &\quad + \frac{1}{D^n} \log \|f_\infty^n[z : w]\| + C_n - \frac{1}{2} \log(|z|^2 + |w|^2). \end{aligned}$$

From one-dimensional theory, we know that $\frac{1}{d^n} \log \|(f_\infty)^n[z : w]\| - \frac{1}{2} \log(|z|^2 + |w|^2)$ converges to a continuous function on L_∞ and $\sum_{p \in E} \lambda_{p,n} \log(|a_p w - b_p z|)$ converges thanks to the previous lemma. The last identity and the fact that $d' < D$ imply the first formula in the proposition. The formula giving m_∞ is then clear by the Poincaré formula. \square

Remark. The previous proof can be applied to all the algebraically stable polynomial maps of \mathbb{C}^2 with indeterminacy points on L_∞ .

The computations in this section are very similar to those in [Dem05]: the author iterates “mappings” in \mathbb{P}^1 of the form $h = [Hp : Hq]$ where H, p , and q are homogeneous polynomials in two variables. A measure μ is introduced and the author proves that it depends continuously on the coefficients of h . Considering $h = [PQ_1, PQ_2]$, we see that here μ is m_∞ , so we deduce that m_∞ depends continuously on the coefficients of P, Q_1 and Q_2 .

2.4. Topological degree.

Let N be a small enough neighborhood of L_∞ and V be a neighborhood of $I(f)$, then there are constants C and C' such that for p in $N \setminus V$, we have:

$$C\|p\|^D \leq \|f(p)\| \leq C'\|p\|^D.$$

Let us assume here that the considered mapping satisfies in addition: for all $I \in I(f)$, there exist a number l_I , a neighborhood $V(I)$ of I , a neighborhood $V(f_\infty(I))$ of $f_\infty(I)$ and constants C_1 and C_2 such that for all $p \in V(I)$ with $f(p) \notin V(f_\infty(I))$, we have:

$$(2) \quad C_1\|p\|^{l_I} \leq \|f(p)\| \leq C_2\|p\|^{l_I}.$$

This condition is easy to check in practice. Under these assumptions, we can compute the topological degree of f which is the mass of the pull-back of any probability measure by f . The difference with the case with no dynamics on L_∞ is that we have to count the number of preimages of a generic line by f_∞ . We have the following proposition:

Proposition 2.9. *Let $f \in \mathcal{G}$ satisfying (2). Then the topological degree of f is given by:*

$$d_t = \sum_{I \in I(f)} l_I \alpha_I + d' D.$$

In particular, we have $d_t > D$.

Proof: Let L be a generic line, we consider the probability measure $[L_\infty] \wedge [L]$ (which is the Dirac mass at the intersection of L and L_∞). By definition, its pull back by f is of mass d_t . After some change of coordinates, we can assume that the point $[1 : 0 : 0]$ is not on L and $f^{-1}(L)$. So we work in the coordinates $(u, v) = (Z/W, T/W)$ where a potential of $L_\infty = (v = 0)$ is $\varphi(u, v) = \log |v|$. We must compute:

$$\int_{\mathbb{P}^2} f^*([L_\infty] \wedge [L]) = \int_{f^{-1}(L)} dd^c(\varphi \circ f).$$

For each I in $I(f)$, let \mathbb{B}_I be a bidisk in $V(I)$ for the (u, v) coordinates, and for each p in $f_\infty^{-1}(L_\infty \cap L)$ let \mathbb{B}_p be a bidisk around p . Since L is a

generic line, we can assume that $f_\infty^{-1}(L_\infty \cap L) \cap I(f) = \emptyset$ and that all those bidisks are disjoint. The previous integral becomes:

$$d_t = \sum_{I \in I(f)} \int_{f^{-1}(L) \cap \mathbb{B}_I} dd^c(\varphi \circ f) + \sum_{p \in f_\infty^{-1}(L_\infty \cap L)} \int_{f^{-1}(L) \cap \mathbb{B}_p} dd^c(\varphi \circ f).$$

Observe that $\varphi \circ f - l_I \log |v|$ is a bounded pluriharmonic function on $\mathbb{B}_I \setminus L_\infty$ thanks to (2), so it defines in fact a pluriharmonic function on \mathbb{B}_I . Hence, on these bidisks, $dd^c(\varphi \circ f)$ is equal to l_I times the current of integration on L_∞ . Using the same argument for \mathbb{B}_p , we deduce:

$$d_t = \sum_{I \in I(f)} \int_{f^{-1}(L) \cap \mathbb{B}_I} l_I dd^c(\log |v|) + \sum_{p \in f_\infty^{-1}(L_\infty \cap L)} \int_{f^{-1}(L) \cap \mathbb{B}_p} Ddd^c(\log |v|)$$

which is what we wanted since $\int_{f^{-1}(L) \cap \mathbb{B}_I} dd^c(\log |v|) = \alpha_I$ is the intersection multiplicity at I of L_∞ and $f^{-1}(L)$ and since there are d' preimages of $L \cap L_\infty$ by f_∞ . \square

3. Structure of the Julia set and of the Green current near infinity

Throughout this section we make the following assumptions:

- f_∞ is uniformly hyperbolic (i.e. the forward orbit of each critical point converges towards some attracting periodic orbit).
- $I(f) \cap J_\infty = \emptyset$.
- The indeterminacy points of f are not periodic under f_∞ .

After a unitary change of coordinates, we can also assume that $[1 : 0 : 0]$ is not in $J_\infty \cup E$. Hence $(u, v) = (Z/W, T/W)$ is a coordinate system of a neighborhood of $L_\infty \setminus [1 : 0 : 0]$ where $L_\infty = (v = 0)$.

We construct suitable boxes (polydisks) around the elements of E such that f defines horizontal-like maps between these boxes.

After decomposing the Julia set into pieces near infinity, we construct a subshift on $E^\mathbb{N}$. We then decompose the Green current along these pieces by pulling-back a smooth vertical positive closed $(1, 1)$ -form in the boxes which gives the Green current in a neighborhood of infinity. Observe that the set of maps we consider contains an open set in the space of parameters.

Next, we give an application for the escape rate of f and we explain how to obtain a weaker decomposition in the more general case where some indeterminacy points are in J_∞ . Finally we explain our results through examples.

3.1. Construction of the boxes.

The purpose of this section is to prove the following proposition:

Proposition 3.1. *For all p in E , there is a bidisk Δ_p centered at p such that f induces by restriction a horizontal-like map from Δ_p to Δ_q for all $q \in E$ if $p \in I(f)$ and for $q = f_\infty(p)$ if p is not an indeterminacy point. We denote by $f_{p,q}$ this restriction.*

The bidisks can be taken arbitrarily small. We can choose them so that for all $I \in I(f)$ and all $q \in E - \{I(f)\}$ then $\Delta_I \cap \Delta_q = \emptyset$, and for all $p \in E$ and all $q, q' \in f_\infty^{-1}(p)$ then $\Delta_q \cap \Delta_{q'} = \emptyset$.

Since f_∞ is uniformly hyperbolic, we can put a smooth conformal metric g on L_∞ such that $\|Df_\infty(z)\|_g \geq \lambda > 1$ on J_∞ . Let us remark that E is discrete in the Fatou set $F_\infty := L_\infty \setminus J_\infty$ since the only components of F_∞ are basins of attraction of periodic cycles and $\bar{E} = E \cup J_\infty$ (see [Mil99]). The idea is first to construct disks on L_∞ which will be thickened to get bidisks. So, we use the following lemma:

Lemma 3.2. *There is a constant $c > 0$ such that for all p in E , there exists a disk \mathbb{D}_p for the metric g such that if $p \in I(f)$ and $q \in E$ then $\text{dist}_g(f_\infty(\partial\mathbb{D}_p), \mathbb{D}_q) \geq c$ and if $p \in E \setminus I(f)$ then $\text{dist}_g(f_\infty(\partial\mathbb{D}_p), \mathbb{D}_{f_\infty(p)}) \geq c$. Furthermore, we can choose the radii of those disks to be bounded and arbitrarily small.*

Proof: Let U be an open neighborhood of J_∞ in L_∞ with smooth boundary such that $\|Df_\infty(z)\|_g \geq \rho > 1$ on U and $f_\infty^{-1}U \subset U$. There is only a finite number of elements of E in $L_\infty \setminus U$. Modifying U if necessary, we can assume that $I(f) \cap U = \emptyset$ and $\partial U \cap E = \emptyset$.

For $I \in I(f)$ such that $f_\infty(I) \notin E$, we consider \mathbb{D}_I a disk centered at I on L_∞ for the metric g with $f_\infty(\mathbb{D}_I)$ far from the other points of E . Restricting \mathbb{D}_I if necessary, we can assume that for all p in $f_\infty^{-1}\{I\}$ there is a disk \mathbb{D}_p centered in p on L_∞ such that $f_\infty(\partial\mathbb{D}_p) \cap \mathbb{D}_I = \emptyset$ (we use the fact that f_∞ is open). We iterate this construction with the preimages of all the p till all of them are in U . Of course, we may have to shrink \mathbb{D}_I at each step. We apply this process to all the elements of $I(f)$ such that $f_\infty(I) \notin E$.

Since we assumed the elements of $I(f)$ are not periodic for f_∞ , we then have disks \mathbb{D}_p for all the p in $E \setminus U$ such that $f_\infty(\partial\mathbb{D}_p) \cap \mathbb{D}_{f_\infty(p)} = \emptyset$. Let r be the smallest radius of all these disks. It can be chosen arbitrarily small.

Next, by hyperbolicity, there is some $\varepsilon_0 > 0$ such that f_∞ is injective on any disk $\mathbb{D}_g(z, \varepsilon_0)$ for all z in $f_\infty^{-1}U$, and is close to its differential. Namely, for all $\varepsilon \leq \varepsilon_0$, there is a $\rho' > 1$ such that we have $\mathbb{D}_g(f(z), \rho'\varepsilon) \subseteq$

$f_\infty(\mathbb{D}_g(z, \varepsilon))$ for z in $f_\infty^{-1}U$. Then, for r small enough, we have some r' such that for all p in $E \cap f_\infty^{-1}U$, the disk $\mathbb{D}_p = \mathbb{D}_g(p, r')$ satisfies $f_\infty(\partial\mathbb{D}_p) \cap \mathbb{D}_{f_\infty(p)} = \emptyset$. The existence of the constant $c > 0$ is then clear by construction for $p \in L_\infty \setminus f_\infty^{-1}(U)$ and by hyperbolicity for $p \in E \cap f_\infty^{-1}(U)$. \square

Proof of Proposition 3.1: Recall that the line at infinity is f -attracting: there is a constant $C > 1$ such that for $M = (z, w)$ in \mathbb{C}^2 with $\|M\| \geq A$, we have $\|f(M)\| \geq C\|M\|$ where $\|\star\|$ denotes the euclidean norm. Furthermore:

$$\|M\|^2 = |z|^2 + |w|^2 = \left| \frac{u}{v} \right|^2 + \left| \frac{1}{v} \right|^2.$$

If $p = (u_p, v_p)$, define $\Delta_p = \mathbb{D}_p \times \left\{ |v| < \frac{\epsilon}{\sqrt{1+|u_p|^2}} \right\}$ with ϵ small. For $M = (u, v) \in \Delta_p$, we have that

$$(1 + \nu)^{-1}(|u_p|^2 + 1) \left| \frac{1}{v} \right|^2 \leq \|M\|^2 \leq (1 + \nu)(|u_p|^2 + 1) \left| \frac{1}{v} \right|^2$$

where $\nu > 0$ depends only on the radius of \mathbb{D}_p (since u is uniformly bounded) and goes to zero with it. We take the radii of the \mathbb{D}_p small enough so that the square of the norm of an element in Δ_p is close to $(|u_p|^2 + 1) \left| \frac{1}{v} \right|^2$.

We choose ϵ so that all the Δ_p are in the domain where the infinity is attracting. Restricting r' which is the supremum of the radii of the disks \mathbb{D}_p if necessary, we can assume that $f(\Delta_p) \cap \partial\Delta_q \subset \partial_v\Delta_q$ for all q if p is an indeterminacy point and for $q = f(p)$ otherwise. Now, using the uniform continuity of f in the complement of some small neighborhood of the indeterminacy set and the existence of c in Lemma 3.2, we can choose ϵ so that $f(\partial_v\Delta_p) \cap \Delta_q = \emptyset$ for all $q \in E$ if $p \in I(f)$ and for $q = f(p)$ otherwise. Finally, since the image of any small neighborhood of an indeterminacy point by f contains the whole line at infinity, we have $f(\Delta_p) \cap \Delta_q \neq \emptyset$. The last part of the proposition is clear for r' small enough (we use the hyperbolicity of f_∞ once again here). \square

3.2. Construction of the subshift.

Now, we define the symbolic dynamics which will appear in the decomposition of the Green current. First, we will need to know the degree of the horizontal-like maps $(f_{p,q})$. Recall that α_i is the multiplicity of the indeterminacy point $I_i \in I(f)$. We take the notations of (1). The following lemma is clear:

- Lemma 3.3.** 1. If p is in $E \setminus I(f)$, then the degree of $f_{p,q}$ is the local degree of f_∞ at p ,
 2. if $p = I_j$ is in $I(f)$ and $q \neq f_\infty(I_j)$, then the degree of $f_{p,q}$ is α_j ,
 3. if $p = I_j$ is in $I(f)$ and $q = f_\infty(I_j)$, then the degree of $f_{I_j, f_\infty(I_j)}$ is the sum of α_j and the local degree of f_∞ at I_j .

Define $\Sigma' = E^\mathbb{N}$ and $\Sigma = \{(\beta_n) \in \Sigma', f_{\beta_n, \beta_{n+1}} \text{ exists}\}$, the space of itineraries between indeterminacy points and their preimages. We consider the left shift σ on Σ and Σ' . Define $N = \bigcup_{p \in E} \Delta_p$. For $\beta \in \Sigma$, let us consider:

$$\mathcal{K}_\beta = \{p \in N, f^j(p) \in \Delta_\beta\}.$$

Then, for all $\beta \in \Sigma$, $\overline{\mathcal{K}_\beta}$ is not empty as a decreasing intersection of vertical closed sets in $\Delta_{\beta(0)}$. Let \mathcal{K} be the union of all the \mathcal{K}_β so that $\mathcal{K} \subset N$.

Observe that $T \wedge [L_\infty]$ is the slice of T by $(v = 0)$. Using the formula giving the trace of T on L_∞ and the invariance of T ($f^*T = DT$), we have that:

$$\forall p \in E, \lambda_p = \frac{1}{D} \sum_{q \in E} d_{p,q} \lambda_q$$

with the convention that $d_{p,q} = 0$ if $f_{p,q}$ is not defined. For all $p \in E$, we deduce:

$$(3) \quad 1 = \sum_{q \in E} \frac{d_{p,q} \lambda_q}{D \lambda_p}.$$

For example, if p is not an indeterminacy point, we have that all the $d_{p,q}$ are zero except for $q = f(p)$ and the formula becomes:

$$1 = \frac{d_{p, f(p)} \lambda_{f(p)}}{D \lambda_p}.$$

And if $p = I$ is in the indeterminacy set with $d_{I,q}$ constant (i.e. $f_\infty(I) \notin E$), then:

$$1 = \sum_{q \in E} \lambda_q.$$

Let $A := (a_p^q)_{p,q \in E}$ be the infinite matrix defined by $a_p^q = \frac{d_{p,q} \lambda_q}{D \lambda_p}$. The entry a_p^q can be seen as the *probability to go from Δ_p to Δ_q by f* in term of slice mass (see the proof of Theorem 3.5). Of course, if p is not an indeterminacy point, one always goes to $\Delta_{f(p)}$ (the probability is 1). We

put on Σ the Borel measure ν defined by:

$$\begin{aligned} \nu(\{\beta \in \Sigma, \beta(0) = \beta_0, \dots, \beta(n) = \beta_n\}) &= \lambda_{\beta_0} \times \prod_{i=0}^{n-1} a_{\beta_i}^{\beta_{i+1}} \\ &= \lambda_{\beta_n} \times \prod_{i=0}^{n-1} \frac{d_{\beta_i, \beta_{i+1}}}{D}. \end{aligned}$$

Proposition 3.4. *The left shift σ on Σ defines a subshift for which the measure ν is invariant and mixing.*

Proof: Definitions and facts on symbolic dynamics and especially subshift can be found in [KH95, pp. 156–158], although the authors do not mention subshift on a countable alphabet, all the facts stated there easily generalized to that case apart for the mixing (which is in general false for subshift on a countable alphabet). By (3), we already have for all $p \in E$ that

$$(4) \quad \sum_q a_p^q = 1.$$

What remains to be proved is that the vector (λ_p) is an eigenvector for the matrix ${}^t A$ associated with the eigenvalue 1 (that gives the invariance of ν). That is:

$$(5) \quad \sum_p a_p^q \lambda_p = \lambda_q$$

which is clear again by (3).

Furthermore, the matrix A is transitive in the sense that for each (p, q) the entry of index (p, q) in A^n is strictly positive for some n (it is clear if p is in the indeterminacy set for $n = 1$ and if $p \in f^{-j}(I(f))$, then it is true for $n = j + 1$).

Now we consider only a finite part E_N of E containing the indeterminacy points and their preimages up to some order and regroup the rest of the elements of E in a same box. That way, we get a finite Markov chain, but we lose some part of the information, the application $E \mapsto E_N$ induces an application from the two subshifts which preserves the measure. That gives the mixing since we only need to consider cylinders for the mixing, and a cylinder can be viewed as an element in $(E_N)^{\mathbb{N}}$ if E_N is large enough, and we know finite subshift are mixing. \square

3.3. Decomposition of the Green current.

Let us denote by $\mathcal{L}_{p,q}$ the operator $\frac{1}{d_{p,q}} f_{p,q}^*$ acting on vertical currents. Recall that f is a polynomial map of \mathbb{C}^2 having indeterminacy points on L_∞ which is f -attracting. The map f_∞ is hyperbolic and the indeterminacy points of f are on the Fatou set of f_∞ and not periodic. We can now prove our main theorem:

Theorem 3.5. 1. *There exists an at most countable set $\Theta \subset \Sigma$ such that for all $\beta \in \Sigma \setminus \Theta$, there is a unique current T_β satisfying the following property: for all sequence of currents (S_{k+1}) of bidegree $(1, 1)$, positive, closed, vertical in $\Delta_{\beta(k+1)}$ of slice mass 1, we have:*

$$\mathcal{L}_{\beta(0),\beta(1)} \cdots \mathcal{L}_{\beta(k),\beta(k+1)} S_{k+1} \longrightarrow T_\beta.$$

2. *The Green current T admits the following decomposition in N :*

$$T = \int_{\Sigma} T_\beta d\nu(\beta).$$

Proof: Since $\mathcal{K} = \bigcup \mathcal{K}_\beta$, only a countable number of \mathcal{K}_β have positive Lebesgue measure. Then Theorem 2.3 implies the first part.

For the second part, let S'_p be a smooth positive closed $(1, 1)$ -form in \mathbb{P}^2 such that near L_∞ , S'_p has its support in $\Delta'_p = \mathbb{D}'_p \times \left\{ |v| < \frac{\epsilon}{\sqrt{1+|u_p|^2}} \right\}$, with $\mathbb{D}'_p \Subset \mathbb{D}_p$. Let S_p be the current defined by S'_p in Δ_p , it is a vertical positive closed current. Normalize S'_p so that S_p is of slice mass λ_p . Observe that if $S' = \sum S'_p$ then $\lim \frac{1}{D^n} (f^n)^*(S') = \sum \lim \frac{1}{D^n} (f^n)^*(S'_p) = T$ since $\sum \lambda_p = 1$. Define $S = \sum S_p$. Finally, write:

$$\Sigma_n = \{(a_0, a_1, \dots, a_{n-1}) \in E^n \mid \exists \beta \in \Sigma, \forall i \leq n-1, a_i = \beta(i)\}$$

and for $a \in \Sigma_n$, write C_a for the cylinder:

$$\{\beta \in \Sigma \mid \beta(0) = a_0, \dots, \beta(n-1) = a_{n-1}\}.$$

Pulling back S by f gives:

$$\begin{aligned} \frac{1}{D} f^* S &= \frac{1}{D} \sum_{p,q \in E} f_{p,q}^* S_q \\ &= \sum_{\beta \in \Sigma_1} \frac{d_{\beta(0),\beta(1)}}{D} \mathcal{L}_{\beta(0),\beta(1)} S_{\beta(1)}. \end{aligned}$$

The bidisks Δ_p are not all disjoint, still, for a given p , the bidisks corresponding to the preimages of p and to the points of indetermination

are disjoint, so we can write uniquely $f^* S_p = \sum_{q \in f^{-1}(p) \cup I(f)} S_q''$ where $S_q'' = f_{q,p}^* S_p$ is a vertical positive closed current in Δ_q . We iterate:

$$\begin{aligned} \frac{1}{D^k} (f^k)^* S &= \sum_{\beta \in \Sigma_k} \frac{\prod_{i=0}^{k-1} d_{\beta(i), \beta(i+1)}}{D^k} \mathcal{L}_{\beta(0), \beta(1)} \cdots \mathcal{L}_{\beta(k-1), \beta(k)} S_{\beta(k)} \\ &= \sum_{\beta \in \Sigma_k} \nu(C_\beta) \mathcal{L}_{\beta(0), \beta(1)} \cdots \mathcal{L}_{\beta(k-1), \beta(k)} \frac{S_{\beta(k)}}{\lambda_k}. \end{aligned}$$

The left hand side goes to the Green current T . By the first part of the theorem, the general term of the right hand side tends to T_β for β generic so we get the result by dominated convergence. \square

Remark. The dynamics of f near infinity is semi-conjugated to the sub-shift σ in the sense that $f(\mathcal{K}_\beta) \subset \mathcal{K}_{\sigma(\beta)}$.

We see that the current gives full mass to \mathcal{K} which does not meet J_∞ . So, as announced in the introduction, the local stable manifolds to the points of the Julia set of f_∞ do not carry any part of the Green current, but they are contained in its support.

3.4. Escape rate.

We take $f \in \mathcal{G}$ satisfying the condition (2), we also suppose that $f_\infty(I(f)) \cap E = \emptyset$ (else, we would have p in E such that $f(p) \in f(V(I(f)))$). We want to compute the possible values of the *upper escape rate* \bar{l} where $\log(\bar{l}) = \limsup \frac{1}{n} \log^+ \log^+ \|f^n\|$ which becomes $\limsup \frac{1}{n} \log \log(\|f^n\|)$ in N . In the same way, we define the *lower escape rate* \underline{l} and we are interested in knowing where these two functions match up, in which case we note l their common value which we simply call the *escape rate*.

For $p \in E \setminus I(f)$, we set $l_p = D$. We have the following lemma:

Lemma 3.6. *Let $\beta \in \Sigma$ and $q \in \mathcal{K}_\beta$. We have:*

$$\frac{1}{n} \log \log \|f^n(q)\| = \frac{1}{n} \log(l_{\beta(0)} l_{\beta(1)} \cdots l_{\beta(n-1)}) + O\left(\frac{\log n}{n}\right).$$

Proof: We have constants c_1 and c_2 such that:

$$c_1 \leq \log \|f^{j+1}(q)\| - l_{\beta(j)} \log \|f^j(q)\| \leq c_2.$$

Taking a combination of these inequalities for $j \leq n - 1$ gives:

$$c_1 \left(\sum_{j=0}^{n-1} l_{\beta(j+1)} \cdots l_{\beta(n-1)} \right) + l_{\beta(0)} \cdots l_{\beta(n-1)} \log \|q\| \leq \log \|f^n(q)\|,$$

with a similar inequality for the right hand side. Taking the logarithm and dividing by n give:

$$\begin{aligned} & \left| \frac{1}{n} \log \log \|f^n(q)\| - \frac{1}{n} \log(l_{\beta(0)} \dots l_{\beta(n-1)}) \right| \\ & \leq \frac{1}{n} \log \left(\log \|q\| + C \sum_{j=0}^{n-1} \frac{1}{l_{\beta(0)} \dots l_{\beta(j-1)}} \right). \end{aligned}$$

The sum in the right hand side is a $O(n)$ which concludes the proof. \square

Choosing a suitable β , we deduce from the lemma that the range of the escape rate in N is $[\min l_I, D]$ (the details are left to the reader). In this case, it is interesting to observe that the set of possible escape rates is an interval which is a new property for polynomial mappings. Let λ denote the slice mass $1 - \sum_{I \in I(f)} \lambda_I$ of T outside a neighborhood of $I(f)$. We have the following theorem:

Theorem 3.7. *For $\|T\|$ -almost every point q in N , the escape rate $l(q)$ exists and is equal to $D^\lambda \prod_{I \in I(f)} l_I^{\lambda_I}$.*

Proof: Since the left shift σ is ergodic for ν , the Birkhoff's ergodic theorem yields that for ν -almost every β :

$$\exp \left(\frac{1}{n} \sum_{i=0}^{n-1} \log l_{\sigma^i(\beta)(0)} \right) \longrightarrow \exp \left(\int_{\Sigma} \log l_{\beta(0)} d\nu \right) = D^\lambda \prod_{I \in I(f)} l_I^{\lambda_I}.$$

And the theorem follows from the previous lemma and Theorem 3.5. \square

3.5. Generalization.

In the case where some indeterminacy points are on J_∞ (possibly periodic), we can obtain a decomposition of the Green current by building a cover of J_∞ by disks such that for all \mathbb{D} in this cover, there exist disjoint disks $\mathbb{D}_1, \mathbb{D}_2, \dots, \mathbb{D}_{d'}$ in the cover with $f_\infty^{-1}(\mathbb{D}) \subset \mathbb{D}_1 \cup \mathbb{D}_2 \cup \dots \cup \mathbb{D}_{d'}$ and $\mathbb{D} \Subset f_\infty(\mathbb{D}_i)$ for all $i \leq d'$. The trick is to have two disks \mathbb{D}_I and \mathbb{D}'_I around each indeterminacy point $I \in I(f)$ so that $\partial f_\infty(\mathbb{D}_I) \cap \overline{\mathbb{D}} = \emptyset$ or $\partial f_\infty(\mathbb{D}'_I) \cap \overline{\mathbb{D}} = \emptyset$. Finally, we follow the construction of Section 3.1 with U being replaced by the union of all those disks.

This time we only have a finite number of bidisks and when we pull back the Green current near some point of E to an indeterminacy point I in J_∞ , we may have to choose between the two bidisks centered at I in order to have a horizontal-like map. We only get a finite subshift, but taking a finer cover, we get more precision on the decomposition (only

on a smaller neighborhood of L_∞). Somehow the decomposition is not intrinsic because we do not pull back according to the itinerary but it assures that the Green current is not extremal in a neighborhood of L_∞ .

3.6. Examples.

First let us explain our results in two examples where the dynamics at infinity is linear.

Example 1. Consider the case where f_∞ is given by $u \mapsto 2u$ and where the indeterminacy set is reduced to $(1, 0)$ with multiplicity 1 in the (u, v) coordinates (thanks to Proposition 2.5, we know this case exists, take for example $f(z, w) = C(2z(z - w) + z, w(z - w))$ for C large enough). Then, using Proposition 2.8, we find that:

- $E = \{p_n = (\frac{1}{2^n}, 0), n \geq 0\}$.
- $\lambda_n = \frac{1}{2^n}$.
- The matrix of the subshift is:

$$\begin{pmatrix} \frac{1}{2} & \frac{1}{4} & \frac{1}{8} & \cdots \\ 1 & 0 & 0 & \cdots \\ 0 & 1 & 0 & \cdots \\ \vdots & \vdots & \vdots & \ddots \end{pmatrix}.$$

An element $\beta \in \Sigma$ can be written $(p_{n_1}, p_{n_1-1}, \dots, p_0, p_{n_2}, \dots, p_0, \dots)$ for some sequence (n_i) in \mathbb{N} . The dynamics in the space of itineraries is simple: a point in \mathcal{K}_β where $\beta_0 = p_{n_1}$ is sent near p_{n_1-1} then near p_{n_1-2} and so on until it arrives near p_0 , in which case it can be sent near any element of E since p_0 is an indeterminacy point.

Example 2. This time, we still take f_∞ given by $u \mapsto 2u$ and we suppose that the indeterminacy points are $I_0 = (2, 0)$ and $I_1 = (1, 0)$ with multiplicity 1 in the (u, v) coordinates, so $D = 3$ (for example: $f(z, w) = (2z(z - w)(z - 2w) + z^2, w(z - w)(z - 2w))$). In this case, we have that $f_\infty^{-1}I_0 = I_1$. Again, using Proposition 2.8, we find that:

- $E = \{p_n = (\frac{1}{2^{n-1}}, 0), n \geq 0\}$.
- We have $\lambda_0 = \lambda_{I_0} = \frac{1}{3}$, $\lambda_1 = \lambda_{I_1} = \frac{4}{9}$, $\lambda_{p_n} = \frac{4}{3^{n+1}}$.
- The matrix of the subshift is:

$$\begin{pmatrix} \frac{1}{3} & \frac{4}{9} & \frac{4}{27} & \frac{4}{3^4} & \cdots \\ \frac{1}{2} & \frac{1}{3} & \frac{1}{9} & \frac{1}{3^3} & \cdots \\ 0 & 1 & 0 & 0 & \cdots \\ 0 & 0 & 1 & 0 & \cdots \\ \vdots & \vdots & \vdots & \vdots & \ddots \end{pmatrix}.$$

The interesting fact here is that the entries of the second row are not proportionnal to the the slice mass, indeed a point near I_1 will have “more chances” to be sent on Δ_{I_0} by f since $f_\infty(I_1) = I_0$.

Example 3. Now, we consider the case were the indeterminacy points are in the exceptionnal set of f_∞ (namely $f^{-1}(I) = I$). Observe that this case does not meet the hypothesis of Theorem 3.5 since the indeterminacy points are periodic. For example, let $f : (z, w) \mapsto (z^3 + w^2, zw^2)$.

By Proposition 2.5, L_∞ is f -attracting. We even have $\|f(z, w)\| \geq \|(z, w)\|^2$ for $\|(z, w)\|$ large enough. The meromorphic extension of f to \mathbb{P}^2 is given by: $f([Z : W : T]) = [Z^3 + TW^2 : ZW^2 : T^3]$. The indeterminacy set of f is reduced to $I_0 = [0 : 1 : 0]$ and the dynamics at infinity is given by $f_\infty : [z : w : 0] \mapsto [z^2 : w^2 : 0]$ (so $f_\infty^{-1}(I_0) = I_0$). Thus f is in \mathcal{G} and is algebraically stable.

The topological degree d_t of f , which is by definition the number of preimages of a generic point, is equal to 8 (solve $f(z, w) = (0, 1)$). It is greater than the algebraic degree.

We use the coordinates $(u, v) = (\frac{Z}{W}, \frac{T}{W})$ in which L_∞ is given by $(v = 0)$. The map f becomes:

$$f : (u, v) \mapsto \left(\frac{u^3 + v}{u}, \frac{v^3}{u} \right).$$

In these coordinates, the point I_0 becomes $(0, 0)$. The map f_∞ is given by $u \mapsto u^2$ for which the Julia set J_∞ is the unit circle ($|u| = 1$). We have the following lemma:

Lemma 3.8. *Let $V = \{(u, v), |u| < \frac{1}{2} \text{ and } |v| < \frac{1}{4}|u|^3\}$, then $f(V) \subset V$.*

Proof: Observe that $(0, 0)$ is not in V since f is not defined there. Let (u, v) be in V . We check:

$$\frac{|u^3 + v|}{|u|} \leq |u|^2 + \frac{|v|}{|u|} < \frac{1}{4} + \frac{|u|^2}{4} < \frac{1}{2}.$$

We also have the inequalities:

$$\frac{|u^3 + v|}{|u|} \geq |u|^2 - \frac{|v|}{|u|} > |u|^2 - \frac{|u|^2}{4} > \frac{1}{2}|u|^2$$

$$\frac{|v|^3}{|u|} < \frac{1}{4^3}|u|^8.$$

It is then sufficient to check that:

$$\frac{1}{4^3}|u|^8 < \frac{1}{4} \left(\frac{1}{2}|u|^2 \right)^3$$

which is obvious. \square

We deduce from the lemma that V is in the Fatou set since the sequence of iterates is normal there. Let then $\mathbb{D}_0 \subset \mathbb{D}_1$ be disks on L_∞ centered on I_0 , small enough to be contained in V , with $f_\infty^{-1}(\mathbb{D}_0) \Subset \mathbb{D}_1$. Let \mathbb{D}_2 be a disk centered on $[1 : 0 : 0]$ containing the Julia set of f_∞ with $\partial\mathbb{D}_2 \subset V$. We have that $f^{-1}(\mathbb{D}_2) \Subset \mathbb{D}_2$. We can shrink those disks to have $\mathbb{D}_1 \cap \mathbb{D}_2 = \emptyset$.

As in Proposition 3.1, we want to “thicken” those disks in order to have bidisks such that f defines by restriction horizontal-like maps between them. Close to I , the norm of a point (in the (z, w) coordinates) is given by $|v|^{-1}$, but next to $[1 : 0 : 0]$, it is controlled by $\frac{|u|}{|v|}$ so we use the coordinates $(u', v') = \left(\frac{T}{Z}, \frac{W}{Z}\right)$ there. Then, we define $\Delta_0 = \mathbb{D}_0 \times (|v| < \varepsilon)$, $\Delta_1 = \mathbb{D}_1 \times (|v| < \varepsilon)$ and $\Delta_2 = \mathbb{D}_2 \times (|u'| < \varepsilon')$. Take ε and ε' small enough so that the vertical boundaries of the bidisks are relatively compact in V . Observe that $\Delta_1 \setminus \Delta_0 \subset V$ is in the Fatou set of f .

Recall that since I_0 is an indeterminacy point, any neighborhood of I_0 is sent on the whole L_∞ . Since L_∞ is f -attracting, and by uniform continuity of f away from any neighborhood of I_0 , we can chose ε and ε' small enough so that:

- $f: \Delta_1 \rightarrow \Delta_0$ defines a horizontal-like map of degree 3 denoted by $f_{1,0}$.
- $f: \Delta_1 \rightarrow \Delta_2$ defines a horizontal-like map of degree 1 denoted by $f_{1,2}$.
- $f: \Delta_2 \rightarrow \Delta_2$ defines a horizontal-like map of degree 2 denoted by $f_{2,2}$.

Next, we consider the Green current T of f . We know that its support is contained in the Julia set of f (see [Sib99]). So we know that in some neighborhood of infinity, T can be written as $T_1 + T_2$ where T_1 and T_2 are vertical positive closed currents in $\Delta_0 \subset \Delta_1$ and in Δ_2 . Pulling-back T_1 and T_2 and using the invariance of T , we see that:

$$\frac{1}{3}f^*T = T = T_1 + T_2.$$

So:

$$T_1 = \frac{1}{3}f_{1,0}^*T_1 + \frac{1}{3}f_{1,2}^*T_2$$

$$T_2 = \frac{1}{3}f_{2,2}^*T_2.$$

Calling m_1 and m_2 the slice masses of T_1 and T_2 , we can compute them using the previous equation and the fact that the pull-back of a vertical current of slice mass m by a horizontal-like map of degree d is of slice mass dm . So, we have:

$$m_1 = m_1 + \frac{1}{3}m_2$$

$$m_2 = \frac{2}{3}m_2.$$

Hence, $m_2 = 0$ and so $T_2 = 0$. In particular, the support of the Green current of f is *strictly contained* in the Julia set J since the stable manifolds associated to the Julia set J_∞ of f_∞ are in J but $\text{supp}(T)$ does not meet J_∞ . In [FS95], there is a different example of such phenomenon.

For $\varepsilon > 0$, we consider the small perturbation f_ε defined by:

$$f_\varepsilon : (z, w) \longmapsto ((z + \varepsilon w)z^2 + w^2, (z + \varepsilon w)w^2).$$

We check that f_ε gives the same map at infinity than f and that the indeterminacy point is now $I_\varepsilon = [-\varepsilon : 1 : 0]$. We see that the preimages of I_ε accumulate on the Julia set of $f_{\varepsilon\infty}$ and we have seen that they are on the support of the Green current which contrasts with what happens for f . Here the support of the Green current does not vary continuously with ε whereas the Green current itself varies continuously for the weak topology.

The method used to study that example can be generalized to the case where the indeterminacy set is contained in the exceptional set. For the case where $f_\infty = z^n$ and $I(f) = \{[1 : 0 : 0], [0 : 1 : 0]\}$ we also need to use the method of Section 3.5 (we take two bidisks around each indeterminacy point). The symbolic dynamics that arises here is a finite subshift at two elements.

References

- [BJ00] E. BEDFORD AND M. JONSSON, Dynamics of regular polynomial endomorphisms of \mathbb{C}^k , *Amer. J. Math.* **122**(1) (2000), 153–212.

- [Dem05] L. DEMARCO, Iteration at the boundary of the space of rational maps, *Duke Math. J.* **130**(1) (2005), 169–197.
- [DDS05] T. C. DINH, R. DUJARDIN AND N. SIBONY, On the dynamics near infinity of some polynomial mappings in \mathbb{C}^2 , *Math. Ann.* **333**(4) (2005), 703–739.
- [DS03] T. C. DINH AND N. SIBONY, Dynamique des applications d'allure polynomiale, *J. Math. Pures Appl. (9)* **82**(4) (2003), 367–423.
- [DS04] T. C. DINH AND N. SIBONY, Dynamique des applications polynomiales semi-régulières, *Ark. Mat.* **42**(1) (2004), 61–85.
- [Duj04] R. DUJARDIN, Hénon-like mappings in \mathbb{C}^2 , *Amer. J. Math.* **126**(2) (2004), 439–472.
- [FJ05] C. FAVRE AND M. JONSSON, Eigenvaluations, available at www.arxiv.org/pdf/math.DS/0410417 (2005).
- [FS95] J. E. FORNÆSS AND N. SIBONY, Complex dynamics in higher dimension. II, in: “*Modern methods in complex analysis*” (Princeton, NJ, 1992), Ann. of Math. Stud. **137**, Princeton Univ. Press, Princeton, NJ, 1995, pp. 135–182.
- [KH95] A. KATOK AND B. HASSELBLATT, “*Introduction to the modern theory of dynamical systems*”, With a supplementary chapter by Katok and Leonardo Mendoza, Encyclopedia of Mathematics and its Applications **54**, Cambridge University Press, Cambridge, 1995.
- [Mil99] J. MILNOR, “*Dynamics in one complex variable. Introductory lectures*”, Friedr. Vieweg & Sohn, Braunschweig, 1999.
- [Sib99] N. SIBONY, Dynamique des applications rationnelles de \mathbb{P}^k , in: “*Dynamique et géométrie complexes*” (Lyon, 1997), Panor. Synthèses **8**, Soc. Math. France, Paris, 1999, pp. ix–x, xi–xii, 97–185.

Mathématiques - Bât. 425, UMR 8628
 Université Paris-Sud
 91405 Orsay
 France
 E-mail address: gabriel.vigny@math.u-psud.fr

Primera versió rebuda el 12 de juliol de 2006,
 darrera versió rebuda el 16 de novembre de 2006.